



Contribution ID: 24

Type: **Talk**

## Temporal evolution of magnon-magnon interaction in a magnetic vortex

*Tuesday, 4 June 2019 17:30 (20 minutes)*

Brillouin light scattering microscopy (TR- $\mu$ BLS) measurements on a Permalloy disk magnetized in the vortex state are presented. By applying a homogenous out-of-plane AC field with sufficiently large amplitudes it is possible to drive the spin waves in the nonlinear regime and initiate three- and four-magnon scattering processes.

Time resolved  $\mu$ BLS is used to show that these pumping conditions cause cascades of different types of magnon-magnon interactions. It is shown, that the temporal transition of different scattering mechanisms can be tuned by the excitation frequency and amplitude.

The authors acknowledge financial support from the Deutsche Forschungsgemeinschaft within programme SCHU 2922/1-1.

**Primary author:** Mr HULA, Tobias (Helmholtz-Zentrum Dresden-Rossendorf)

**Co-authors:** Dr SCHULTHEISS, Katrin (Helmholtz-Zentrum Dresden-Rossendorf); Mr KÖRBER, Lukas (Helmholtz-Zentrum Dresden-Rossendorf); Mr HACHE, Toni (Helmholtz-Zentrum Dresden-Rossendorf); Dr SCHULTHEISS, Helmut (Helmholtz-Zentrum Dresden-Rossendorf)

**Presenter:** Mr HULA, Tobias (Helmholtz-Zentrum Dresden-Rossendorf)

**Session Classification:** Spin wave and magnonics

**Track Classification:** Spin wave and magnonics