



Contribution ID: 19

Type: **Talk**

## Stripe discommensuration and spin dynamics in $\text{Pr}_{3/2}\text{Sr}_{1/2}\text{NiO}_4$

*Tuesday, 4 June 2019 18:10 (20 minutes)*

Inelastic neutron scattering studies on the magnetic excitation in a stripe-ordered  $\text{Pr}_{3/2}\text{Sr}_{1/2}\text{NiO}_4$  at 5K reveal that the magnetic incommensurability ( $\epsilon = 0.4$ ) as result of admixing stripe dicommensuration in the checker-board matrix in  $\text{NiO}_2$  plane. A suggested linear spin-wave model accounting 3D-stripe discommensuration with two-fold exchange interactions between  $\text{Ni}^{2+}$  spins, provides a good agreement with the measured spin wave dispersion up to 64 meV, notably, to describe a slight symmetric shift of the broadened peak in the energy range from 35 to 45 meV. Our results indicate that discommensuration model is essential to consolidate in the LSWT calculation to understand the microscopic effect of doped holes to the spin microstructure.

**Primary authors:** Dr DUTTA, Rajesh (Institut für Kristallographie, RWTH Aachen University); Dr MAITY, Avishek (Institute for Physical Chemistry | Georg-August-University of Goettingen)

**Presenter:** Dr DUTTA, Rajesh (Institut für Kristallographie, RWTH Aachen University)

**Session Classification:** Spin wave and magnonics

**Track Classification:** Spin wave and magnonics