



Contribution ID: 6

Type: Talk

## Experimental and theoretical studies of the lattice dynamics in superconducting $\text{BaNi}_2(\text{As}_{1-x}\text{P}_x)_2$ .

Wednesday 16 September 2015 10:15 (25 minutes)

We present a combination of Thermal Diffuse Scattering (TDS) and Inelastic X-ray Scattering (IXS) measurements, and Density Functional Perturbation Theory (DFPT) calculations of the lattice dynamics in superconducting  $\text{BaNi}_2(\text{As}_{1-x}\text{P}_x)_2$ . At  $T_s=130$  K,  $\text{BaNi}_2\text{As}_2$  undergoes a structural phase transition from a tetragonal to a triclinic crystal structure. In addition, superconductivity emerges at  $T_c=0.7$  K. Substitution of the arsenic with phosphorus leads to a suppression of  $T_s$ . Finally, at a substitution level of 7%, the structural phase transition is completely suppressed and the  $T_c$  jumps from 0.7 to 3.3 K. It is believed that the nature of the superconductivity displayed by this material is of the conventional BCS type. Furthermore, specific heat measurements suggest that a “Giant” phonon softening is responsible for the jump in  $T_c$ . In order to determine if such a large phonon softening does occur in  $\text{BaNi}_2(\text{As}_{1-x}\text{P}_x)_2$ , we have undertaken a combined experimental and theoretical study of the lattice dynamics of this material. TDS measurements on  $\text{BaNi}_2\text{As}_2$  have identified locations of significant diffuse scattering. Upon cooling, the scattering from these diffuse regions becomes stronger, collapsing into true Bragg structural reflections below  $T_s$ . The IXS measurements and DFPT calculations identify this to be the location of a significant softening of at least one phonon mode. Our results demonstrate that these phonons are strongly coupled to the structural phase transition, and therefore it is a strong candidate for being responsible for the jump in  $T_c$ . In order to ascertain if this is correct, we have extended our TDS and IXS measurements to phosphorus doped  $\text{BaNi}_2\text{As}_2$ . Results from these latest set of measurements will be presented. In addition, our results will be compared to the lattice dynamics of the iron pnictide unconventional superconductors.

**Author:** Dr FORREST, Thomas (ESRF - The European Synchrotron)

**Co-authors:** Dr BOSAK, Alexei (ESRF - The European Synchrotron); Dr WEHINGER, Bjorn (University of Geneva); Dr ROTUNDU, Costel (Stanford University)

**Presenter:** Dr FORREST, Thomas (ESRF - The European Synchrotron)

**Session Classification:** Excitations of strongly correlated electron systems

**Track Classification:** DyProSo2015 Main track