



Contribution ID: 99

Type: Poster

## High-resolution spectroscopy and diffraction at TRISP

*Wednesday, 9 December 2020 17:40 (20 minutes)*

We present the capabilities of TRISP both for high-resolution spectroscopy and diffraction and show typical experimental examples. TRISP is a thermal three axis spectrometer incorporating the resonant spin-echo technique. Typical applications include the measurement of linewidths of phonons and spin excitations in an energy range 0.5-50meV, and the energy width of quasi-elastic scattering, originating, for example, from critical magnetic fluctuations. Neutron Larmor diffraction (LD) is a high-resolution technique which permits the measurement of lattice spacings  $d_{hkl}$  and their distribution  $\Delta d_{hkl}$ . The latter arises, for example, from micro-strains, magnetostriction, structural and magnetic domains, or from a small splitting of Bragg peaks, resulting from distortions of the crystal lattice. The resolution of Larmor diffraction at TRISP is  $10^{-6}$  (relative) for the lattice spacing and one order of magnitude less for the distribution width.

**Primary author:** Dr KELLER THOMAS (MPI outstation at the FRM II)

**Co-authors:** Mr TRALMER, Franz; Prof. KEIMER, Bernhard

**Presenter:** Dr KELLER THOMAS (MPI outstation at the FRM II)

**Session Classification:** Joint poster session of MLZ User Meeting and DN2020

**Track Classification:** DN: Instrumentation