



Contribution ID: 511

Type: Poster

Performing an accurate measurement of Δb_i of ^3He using NSE

Tuesday 21 March 2023 16:00 (2 hours)

We recently performed measurements of the neutron incoherent scattering lengths for ^{129}Xe and ^{131}Xe . As a part of those measurements the Δb_i of ^3He was also measured for a short time, only 6 hours, as a calibration. The results of this measurement indicate a very precise value could be obtained for a typical length experiment on NSE. Theories which take the data from measurements in three-body nuclear systems and predict what should be seen in a 4-body system like ^4He or $n\text{-}^3\text{He}$, which is an essential step needed to check the internal consistency of the three-body force extraction from systems with only three bodies, do not quite agree with the data by amounts too large to be explained only by 4-body forces. Furthermore, in the specific case of the $n\text{-}^3\text{He}$ system, the two best measurements of the $n\text{-}^3\text{He}$ incoherent scattering length differ by 3σ . During the course of our test measurement we found several important experimental and theoretical factors relevant to a measurement with precision beyond the error of the current measurements. We estimate that we could obtain a 1% measurement for b_i of ^3He in one week.

Authors: Prof. GOODSON, Boyd (Southern Illinois University); BABCOCK, Earl; Mr LU, Hao (Indiana University); SNOW, W. Michael (Indiana University Bloomington)

Presenter: BABCOCK, Earl

Session Classification: Poster session TUESDAY

Track Classification: Fundamental Science