



Contribution ID: 33

Type: Poster

Magnetic excitations in the zigzag-chain compound $\text{KCu}_4\text{P}_3\text{O}_{12}$

Monday 20 March 2023 16:00 (2 hours)

We report on the results of inelastic neutron scattering (INS) of $\text{KCu}_4\text{P}_3\text{O}_{12}$. Cu^{2+} ions carry $S = 1/2$ spins. Four types of nearest-neighbor exchange interactions form the eight-spin zigzag-chain ($J_1 - J_2 - J_3 - J_4 - J_3 - J_2 - J_1$ chain). We evaluated the exchange interactions as $J_1 = -8.5$ meV (antiferromagnetic), $J_2 = -2.7$ meV, $J_3 = -3.9$ meV, and $J_4 = 6.2$ meV (ferromagnetic) from magnetic susceptibility and magnetization curves at several temperatures using a data-driven technique based on machine learning proposed by Tamura and Hukushima. In the eight-spin zigzag-chain with the values of the exchange interactions, the ground state (GS) is the spin-singlet state. The first-triplet, second-triplet, third-quintet, and fourth-triplet excited states (1ES, 2ES, 3ES, and 4ES, respectively) are located at 2.9, 4.2, 6.7, and 8.5 meV, respectively.

We performed INS experiments on $\text{KCu}_4\text{P}_3\text{O}_{12}$ powder using the HRC spectrometer at J-PARC (Proposal ID 2020B0026). We used incident neutrons with the energy 15.3 and 25.4 meV. Excitations at 5.5 K are most apparent at around $\omega = 3$ meV and $Q = 0.7 \text{ \AA}^{-1}$. The 3 meV excitation exists in a wide Q range, indicating cluster excitation, and corresponds to the excitation from GS to 1ES (2.9 meV). Excitations are seen up to about 6 meV, indicating the existence of the excitation from GS to 2ES (4.2 meV). We can see weak excitations around 9 meV, corresponding to the excitation from GS to 4ES (8.5 meV). Probably, we also observed the excitation from 1ES to 3ES (3.8 meV) at high-temperature data. As a result, the calculated energies of 1ES, 2ES, 3ES, and 4ES from GS are consistent with the experimental results. We will discuss the Q dependence of the INS intensity.

Author: MATSUO, Masashi (National Institute for Materials Science)

Co-authors: Prof. HUKUSHIMA, Koji (The University of Tokyo); Dr TAMURA, Ryo (National Institute for Materials Science); Dr ASAI, Shinichiro (The University of Tokyo); Prof. MASUDA, Takatsugu (The University of Tokyo)

Presenter: MATSUO, Masashi (National Institute for Materials Science)

Session Classification: Poster Session MONDAY

Track Classification: Magnetism, Superconductivity, Topological Systems, Magnetic Thin Films and other electronic phenomena