



Model cellular membranes: From flat to strongly curved structures

Tuesday 21 March 2023 16:00 (2 hours)

Cellular membranes are essential components of cells that compartmentalize cellular events, control communication between compartments and with the exterior, enable the formation of gradients of ions and other solutes, and provide a proper environment for the large percentage of proteins that are membrane bound or associated to it¹. The ability to carry out this diverse array of essential functions requires a high degree of dynamic organization at the nanoscale. It is proposed that the different functions are compartmentalized into different domains by lipid-lipid and/or lipid-protein interactions, but studying these systems have proved difficult due to the nature of cellular membranes and the requirement of sophisticated techniques to analyze them at the molecular level. The current state of the art on membrane organizations heavily focusses on planar membranes² despite these membranes often being highly curved as in dynamic processes such as invagination and vesiculation, but also in organelles such as the endoplasmic reticulum and the mitochondria. Neutron scattering combined with deuteration is an ideal technique to study the structure and dynamics of multicomponent systems where different parts of the system can be highlighted individually^{3–5}. For example, Neutron Reflection (NR) allows to extract depth profiles of rather complex biointerfaces² and revealed the overall structure of lipid bilayers as a function of composition^{6,7}, the dependency of the core thickness on acyl chain type⁷, and the position of cholesterol in the bilayer^{8,9}, the flip/flop¹⁰, and the ability of membranes to exchange lipids across bilayers³, among other parameters such as fluctuations on floating bilayers¹¹. Similar depth profile information can be extracted for lipid vesicles from small angle neutron scattering (SANS) but the accuracy is lower as compared to NR due to the smearing by the inherent orientational averaging and lower dQ/Q . For example, using SANS the membrane thickness of photolipid vesicles was shown to be tuned in response to illumination by UV and blue light¹². Both NR and SANS lack the possibility to extract lateral correlations, which is accessible to GISANS, such as local adhesion points, clustering of adhesion anchors¹³ as well as lipid rafts¹⁴.

In this talk I will present a range of examples that demonstrate the power of neutron scattering and selective deuteration for studying the structure and composition of model cellular membranes. I will then finally present a new approach based on diffracting scaffolds to study membrane structure as a function of curvature suited for a range of surface sensitive techniques. I will also show fluorescence microscopy data for lipid diffusion on curved membranes among others.

References:

1. Pomorski, T. G., Nylander, T. & Cárdenas, M. Model cell membranes: discerning lipid and protein contributions in shaping the cell. *Adv. Colloid Interface Sci.* 205, 207–220 (2014).
2. Clifton, L. A. et al. Design and use of model membranes to study biomolecular interactions using complementary surface-sensitive techniques. *Adv. Colloid Interface Sci.* 277, 102118 (2020).
3. Browning, K. L. et al. Human Lipoproteins at Model Cell Membranes: Effect of Lipoprotein Class on Lipid Exchange. *Scientific Reports* vol. 7 (2017).
4. Maric, S. et al. Time-resolved small-angle neutron scattering as a probe for the dynamics of lipid exchange between human lipoproteins and naturally derived membranes. *Sci. Rep.* 9, 7591 (2019).
5. Wadsäter, M. et al. Monitoring shifts in the conformation equilibrium of the membrane protein cytochrome P450 reductase (POR) in nanodiscs. *J. Biol. Chem.* 287, 34596–34603 (2012).
6. Luchini, A. et al. Peptide discs as precursors of biologically relevant supported lipid bilayers. *J. Colloid Interface Sci.* 585, 376–385 (2021).
7. Åkesson, A. et al. Composition and structure of mixed phospholipid supported bilayers formed by POPC and DPPC. *Soft Matter* 8, 5658–5665 (2012).
8. Waldie, S. et al. Localization of Cholesterol within Supported Lipid Bilayers Made of a Natural Extract of Tailor-Deuterated Phosphatidylcholine. *Langmuir* 34, 472–479 (2018).
9. Waldie, S. et al. The Production of Matchout-Deuterated Cholesterol and the Study of Bilayer-Cholesterol Interactions. *Sci. Rep.* 9, 5118 (2019).

10. Gerelli, Y., Porcar, L., Lombardi, L. & Fragneto, G. Lipid exchange and flip-flop in solid supported bilayers. *Langmuir* 29, 12762–12769 (2013).
11. Rondelli, V., Fragneto, G., Motta, S., Favero, E. D. & Cantù, L. Reflectivity from floating bilayers: can we keep the structural asymmetry? *J. Phys. Conf. Ser.* 340, 012083 (2012).
12. Urban, P. et al. A lipid photoswitch controls fluidity in supported bilayer membranes. *Langmuir* 36, 2629–2634 (2020).
13. Röttgermann, P. J. F. et al. Cell motility on polyethylene glycol block copolymers correlates to fibronectin surface adsorption. *Macromol. Biosci.* 14, 1755–1763 (2014).
14. Levental, I., Levental, K. R. & Heberle, F. A. Lipid Rafts: Controversies Resolved, Mysteries Remain. *Trends Cell Biol.* 30, 341–353 (2020).

Author: Prof. CARDENAS, Marite (Biofisika Institute)

Presenter: Prof. CARDENAS, Marite (Biofisika Institute)

Session Classification: Poster session TUESDAY

Track Classification: Soft Condensed Matter