



Contribution ID: 24

Type: Poster

Relocation of the cold triple axis spectrometer FLEXX to MLZ, Munich: Larmor diffraction and inelastic scattering

Friday 9 December 2022 15:30 (1h 30m)

The cold triple-axis spectrometer (TAS) FLEXX at HZB is a well-designed and upgraded instrument [1-4]. There is a strong wish that this excellent instrument should be preserved for the community after the shutdown of the HZB neutron source. One attractive gap in the present instrumentation suite of MLZ, is the Larmor-diffraction technique (LD) and, as a natural extension, cold neutron resonant spin echo (NRSE). LD permits the exact measurement of lattice constants and their distribution (internal strains, structural distortions or magnetostriction). In addition, spin correlation lengths in antiferromagnets and antiferromagnetic domain sizes of up to 1 μm can be determined. For looking at time-dependent processes one needs the NRSE mode. TAS comes at no extra cost, as is the main backbone of such an instrument.

The instrument will be placed on a cold neutron source. This will allow for a x4 increase in Q resolution, as well as most importantly access to the low Q region, as compared to the existing TRISP@MLZ. Further, new developments are under way to allow for application of magnetic fields at the sample, hitherto not possible. This opens up new vistas in the exploration of materials. A last attractive option is the possibility to combine high magnetic fields together with cold TAS.

- [1] M. Skoulatos et al., NIMA 647, 100 (2011). [↗](#)
- [2] M.D. Le et al., NIMA 729, 220 (2013).
- [3] F. Groitl et al., Rev. Sci. Instrum. 86 025110 (2015).
- [4] K. Habicht et al., EPJ 83, 03007 (2015).

Authors: SKOULATOS, Markos (TUM); HERTWIG, Martin; GEORGII, Robert; KELLER, Thomas (MPI for Solid State Research, Stuttgart)

Presenter: SKOULATOS, Markos (TUM)

Session Classification: Poster Session

Track Classification: Neutron Methods