MLZ User Meeting 2022



Contribution ID: 158

Type: Talk (20 min + 5 min discussion)

The half- to full-Heusler transition in Ni_{1+x}MnSb

Thursday, 8 December 2022 13:40 (25 minutes)

Structural voids are a defining feature of the half-Heusler structure. Using temperature-dependent neutron diffraction, the disordering of these vacancies is followed across the $C1_b-L2_1$ transition in samples of $Ni_{1+x}MnSb$ for various Ni excesses x, demonstrating the second-order nature of this transition. Structure solution on high-resolution room-temperature neutron diffraction as well as X-ray diffraction data yields a vacancy content in excess of 1 - x, the concentration of constitutional vacancies in the ideal model of site occupations. This is mirrored by the increased lattice constant during temperature-dependent diffraction on cooling, implying an annihilation of the initial vacancy excess at elevated temperatures. The reason for this very slow time-scale of vacancy concentration equilibration as compared to typical systems will be discussed, and consequences for the materials properties will be pointed out.

P. Neibecker et al., phys. status solidi B 259, 2100174 (2022)

 Primary author:
 LEITNER, Michael

 Co-author:
 NEIBECKER, Pascal

 Presenter:
 LEITNER, Michael

 Session Classification:
 Structure Research

Track Classification: Structure Research