



Contribution ID: 84

Type: Poster

Deciphering SASE X-ray Pulse Characteristics with β -Variational Autoencoder Networks

Tuesday, 9 April 2024 18:30 (20 minutes)

Free electron lasers (FEL) play an important role across diverse scientific disciplines. Many experiments can benefit from a non-destructive online photon diagnostic of provided X-ray pulses. One method to obtain information about the pulse profile involves analyzing not the X-ray photons directly, but rather the energy distribution of the electrons downstream of a Self-Amplified Spontaneous Emission (SASE) undulator.

In recent times, neural networks have gained widespread recognition as potent analytical tools spanning various scientific domains. Among these, β Variational Autoencoder (β -VAE) networks stand out for their ability to discern key parameters within unlabeled datasets, even when these parameters are unknown beforehand. This study showcases the application of β -VAEs in characterizing SASE X-ray pulses generated by the free electron laser FLASH in Hamburg. Leveraging data from a Transverse Deflecting Structure (TDS), we demonstrate the β -VAE's capacity to identify the SASE strength, a critical parameter, within real-world data from FLASH. This discovery holds promise in improving the accuracy of lasing off references and therefore enhancing the reconstruction of XUV power profiles.

Primary authors: MÖLLER, Felix (Helmholtz-Zentrum Berlin); GOETZKE, Gesa (Gesa); HARTMANN, Gregor (Helmholtz-Zentrum Berlin); DÜSTERER, Stefan (DESY); OTTO, Thorsten (DESY)

Presenter: GOETZKE, Gesa (Gesa)

Session Classification: Posters

Track Classification: MLC