



Contribution ID: 80

Type: Talk (20 min + 5 min discussion)

Insight on the myoglobin interaction with lipid bilayers within sponge phases using polarized neutron reflectometry

Thursday 5 December 2024 15:50 (25 minutes)

We have studied the heme-protein-lipid interaction with lipid bilayers. This is key for understanding the encapsulation of the iron binding heme-protein like myoglobin and phytohemoglobin, in lipid nanoparticles, LNPs. This type of heme-bound iron can be used to treat anaemia instead of iron in organic salts that is conventionally used. Encapsulation is needed to prevent unwanted proteolytic and redox reactions. We have used sponge phase LNPs with diglycerolmonooleate (DGMO), glycerolmonooleate (GMO) and Polysorbate 80 (P80) as well as LNPs where DGMO was partially replaced with dioleoylglycerophosphocholine (DOPC) to form well defined lipid bilayers mimicking the lipid interface inside the sponge phase. To enhance the contrast and reduce the need for additional solvent contrasts we used silicon substrates with a switchable magnetic contrast layer (MCL) during polarised neutron reflectometry (PNR) experiments [8]. These substrates consisted of 10 nm Fe layer capped with 100 nm SiO₂ layer to protect the Fe layer against corrosion and gave excellent response to the spin state of the neutrons. The formed lipid bilayers had a very high coverage of about 90%, which allowed studies of the interaction of the protein with the lipid interface. The results show that myoglobin interacts so strongly with the lipid bilayer that it was mostly removed from the substrate. The presence of DOPC increased the stability of the bilayer so that remains intact with very low amounts of protein attached.

Primary authors: Dr LUCHINI, Alessandra (Dept. of Physics and Geology, Università degli Studi di Perugia, Perugia, Italy); Mr MACHINGAUTA, Marshall (Physical Chemistry, Dept. of Chemistry, Lund University, Sweden); Dr KÖHLER, Sebastian (Physical Chemistry, Lund University, Sweden); Dr GILBERT, Jennifer (Dept of Life Sciences, Division of Chemical Biology, Chalmers University of Technology, Sweden); Mr YAKYMENKO, Ivan (Department of Physics, Chemistry and Biology (IFM), Linköping University); Prof. BIRCH, Jens (Department of Physics, Chemistry and Biology (IFM), Linköping University, Sweden); Prof. JÄRRENDahl, Kenneth (Department of Physics, Chemistry and Biology (IFM), Linköping University, Sweden); Dr COOPER, Joshaniel F. K. (ESS); Dr STENDAHL, Sjoerd (Dept of Physics and Astronomy, Uppsala University); Prof. LANGRIDGE, Sean (ISIS Neutron and Muon Source, Harwell Science and Innovation Campus, Didcot OX110QX, U.K.); Dr KINANE, Christy (ISIS Neutron and Muon Source, Harwell Science and Innovation Campus, Didcot OX110QX, U.K.); Dr VOROBIEV, Alexey (Institut Max von Laue - Paul Langevin (ILL), Grenoble, France); Dr DEVISHVILI, Anton (Institut Max von Laue - Paul Langevin (ILL), Grenoble, France); Prof. HJÖRVARSSON, Björgvin (Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden); Prof. NYLANDER, Tommy (Physical Chemistry, Lund University, Sweden)

Presenter: Prof. NYLANDER, Tommy (Physical Chemistry, Lund University, Sweden)

Session Classification: Soft Matter

Track Classification: Soft Matter